arXiv:hep-th/9412208v1 22 Dec 1994

EFI 94-64

Dirac Monopoles and Witten's Monopole Equations

Peter G.O. Freund
Enrico Fermi Institute and Department of Physics
The University of Chicago, Chicago, IL 60637, USA

Abstract: A simple solution of Witten's monopole equations is given.

Witten [1] has developed a new, elegant and much simplified approach to the Donaldson theory [2] of 4-manifolds. Instead of studying instantons of non-abelian gauge theory, Witten's dual approach is abelian and starts from the monopole equations

$$F_{AB} = \frac{i}{2}(M_A\overline{M}_B + M_B\overline{M}_A) \tag{1a}$$

$$D_{A\dot{A}}M_A = 0, \quad D_{A\dot{A}} = \partial_{A\dot{A}} + iA_{A\dot{A}} \tag{1b}$$

Here both dotted and undotted spinor indices take the values $1, 2$ and are raised and lowered with the corresponding invariant Levi-Cività tensors. F_{AB} are connected with the abelian field strengths $F_{\mu\nu}$ by

$$F_{AB} = \frac{1}{4}(\sigma^{\mu\nu})_{AB}F_{\mu\nu}, \qquad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \tag{2}$$

Witten noted that in flat space the equations (1) admit no L^2 solutions. We wish to point out that they nevertheless admit the following very simple *non*-L^2 solution

$$A_0 = 0, \qquad \vec{A} = \frac{(-y, x, 0)}{2r(r-z)} \tag{3a}$$

$$\begin{pmatrix} M_1 \\ M_2 \end{pmatrix} = \frac{1}{2r\sqrt{r(r-z)}}\begin{pmatrix} x - iy \\ r - z \end{pmatrix}, \tag{3b}$$

so that

$$\begin{pmatrix} \overline{M}_1 \\ \overline{M}_2 \end{pmatrix} = \frac{1}{2r\sqrt{r(r-z)}}\begin{pmatrix} -(r - z) \\ x + iy \end{pmatrix}. \tag{3c}$$

It is straightforward to check that the fields (3) satisfy Witten's monopole equations (1) (we recall that for $m, n = 1, 2, 3$, $(\sigma^{mn})_{AB} = \epsilon^{mnp}\sigma_p\sigma_2$).

Notice that the potentials (3a) correspond to a singular Dirac (not to an 't Hooft-Polyakov) monopole of minimal magnetic charge. We chose to write these potentials with Dirac-string "showing", rather than using sections [4]. The spinor (3b) is also a familiar object. In fact, by construction, the $SU(2)$ gauge transformation

$$\Omega = \sqrt{2}r \begin{pmatrix} M_1 & \overline{M}_1 \\ M_2 & \overline{M}_2 \end{pmatrix}, \tag{4}$$

(with the M's given by eqs. (3b) and (3c)), is such that $M = \frac{1}{\sqrt{2}r}\Omega U$, U being the unit spinor $U^T = (1\ 0)$, so that this gauge transformation rotates [5] a constant $SU(2)$-triplet Higgs field into a hedgehog. The spinor M thus has the property [6] that $M^\dagger \vec{\sigma} M$ is a Coulomb field, which is essentially the first Witten equation (1a) for the solution (3). What has apparently not been appreciated previously, is the fact that this spinor also obeys the the Weyl-Dirac equation $\sigma^\mu D_\mu M = 0$ which is essentially the second Witten equation (1b) for the solution (3).

The simplicity of the solution (3) (non-L^2 though it is), makes it worthwhile to explore the solubility of Witten's equations in other cases.

I wish to thank J.Gauntlett, J. Harvey and E. Witten for useful comments.

This work was supported in part by the grant NSF: PHY-91-23780

References

[1] E. Witten, preprint IASSNS-HEP-94-96 (hep-th/9411102)

[2] S. Donaldson, J. Diff. Geom. **18**, 279 (1983); Topology **29**, 257 (1993)

[3] P.A.M. Dirac, Phys.Rev. **74**, 817 (1948)

[4] T.T. Wu and C.N. Yang, Phys. Rev. **D12**, 3845 (1975)

[5] J. Arafune, P.G.O. Freund and C.J.Goebel, J. Math. Phys. **16**, 433 (1975)

[6] F. Gürsey in *Gauge Theories and Modern Field Theory*, R. Arnowitt and P. Nath, editors, MIT Press, Cambridge, 1976, p. 369